

=====

**PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS' REGULATIONS
(MEDICINAL PREPARATIONS) - 1986**

Read the package insert carefully in its entirety before using this medicine.

ACAMOL[®] TSINUN DAY
Caplets

DRUGS-ABOUT.COM

COMPOSITION

Each caplet contains:

Paracetamol 500 mg

Pseudoephedrine hydrochloride 25 mg

Inactive Ingredients

Microcrystalline cellulose, Opadry 31F31114 green (lactose, hydroxypropyl methyl cellulose, titanium dioxide, macrogol 4000 [polyethylene glycol], FD&C blue #2/ indicocarmine lake, iron oxide yellow, iron oxide black), povidone, sodium starch glycolate (from potato), stearic acid, magnesium stearate, silicone dioxide.

THERAPEUTIC ACTIVITY

The medicine is intended for the symptomatic relief of common cold and nasal congestion associated with fever and pain. Day care medicine.

WHEN SHOULD THE PREPARATION NOT BE USED?

Do not use this medicine if you are breastfeeding.

Do not use this medicine if you are sensitive to any of its ingredients.

Do not use this medicine concomitantly with monoamine oxidase inhibitors (for depression) or within 14 days of discontinuing such treatment.

Do not take this medicine without consulting a doctor before starting treatment

if you are pregnant or if you are suffering, or have suffered in the past, from impaired function of the heart and/or vascular system, the eyes (e.g. glaucoma), the liver, the kidney/urinary tract, the thyroid gland, the prostate gland, diabetes, hypertension, alcoholism, or viral jaundice.

WARNINGS

Do not use this medicine frequently or longer than 10 consecutive days without consulting your doctor - see below.

If you are sensitive to any type of food or medicine, inform your doctor before commencing treatment with this medicine.

Avoid exceeding the recommended dosage while fasting.

If you consume 3 or more alcoholic drinks every day, consult a doctor whether you should take this medicine or another product for relief of pain and for reduction of temperature. Paracetamol may cause liver damage.

Do not take additional fever-reducing and pain-relieving medicines, or medicines for common cold without consulting a doctor or a pharmacist, in order to prevent paracetamol overdose/toxicity.

DRUG INTERACTIONS

If you are taking another drug concomitantly or if you have just finished treatment with another medicine, inform the attending doctor, in order to prevent hazards or lack of efficacy arising from drug interactions.

This is especially important for medicines belonging to the following groups: stimulants of the central nervous system (appetite suppressants), anticoagulants, antidepressants (including MAOI – see above), antihypertensives and medicines for the heart, cough and cold preparations, hepatic enzyme inducing agents e.g. barbiturates or phenytoin (for seizures) or long term treatment with another analgesic and anti-inflammatory medicines.

SIDE EFFECTS

In addition to the desired effect of the medicine, adverse reactions may occur during the course of taking this medicine, for example: dizziness, nervousness or insomnia.

Effects that require special attention

Allergic reactions (rare): stop treatment and refer to your doctor.

In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your doctor immediately.

Adverse Reactions And Drug Interactions In Children

Parents must inform the attending doctor about any side effects, as well as any additional medicine being taken by a child.

See above for details of special side-effects and drug interactions.

RECOMMENDED DOSAGE UNLESS OTHERWISE PRESCRIBED BY THE DOCTOR:

Adults and children over 12 years of age: 1-2 caplets up to 4 times daily.

Do not take more than 8 caplets daily.

Patients over 60 years of age should consult their doctor before taking this medicine because they may be sensitive to these kind of medicines.

Children 6-12 years: ½ -1 caplet up to 4 times daily.

Do not take more than 5 caplets daily.

Do not exceed the recommended dosage.

This medicine is not usually intended for administration to children under 6 years of age.

If fever continues for more than 3 days, the doctor should be consulted. If fever is absent but your condition has not improved within 7 days the doctor should be consulted.

DIRECTIONS FOR USE

Do not chew! Swallow the caplets with a small amount of water.

Avoid Poisoning!

This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, in order to avoid poisoning.

If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.

Do not induce vomiting unless explicitly instructed to do so by a doctor!

Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.

STORAGE

Store this medicine in a cool and dry place.

Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you.

Do not store different medications in the same package.

Drug Registration No.: 120233004300

MANUFACTURER

Teva Pharmaceutical Industries Ltd.

P.O. Box 3190, Petach-Tikva

DRUGS-ABOUT.COM