


CEFACTOR EXTENDED-RELEASE TABLETS USP

DESCRIPTION

Cefaclor USP, the active ingredient in cefaclor extended release tablets, is a semisynthetic cephalosporin antibiotic for oral administration. Cefaclor USP is chemically designated as 3-chloro-7-D-(2-phenylglycinamido)-3-cephem-4-carboxylic acid monohydrate. The Cefaclor Extended-Release Tablet formulation of cefaclor differs pharmacokinetically from the Cefaclor Immediate-Release Capsule formulation of cefaclor (see CLINICAL PHARMACOLOGY). The structural formula is:



$C_{15}H_{14}ClN_3O_4 \cdot H_2O$

M.W. 385.82

Each cefaclor extended-release tablet contains cefaclor monohydrate equivalent to 500 mg (1.36 mmol) anhydrous cefaclor. In addition, each extended release tablet contains the following inactive ingredients: FD&C blue #2 aluminum lake, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, povidone, stearic acid and talc.

USP Dissolution Test Pending.

CLINICAL PHARMACOLOGY

Pharmacokinetics

The cefaclor extended-release tablet formulation of cefaclor is pharmacokinetically different from the cefaclor immediate-release capsule formulation of cefaclor (see Table 1). No direct comparisons with the suspension formulation of cefaclor have been conducted; therefore, there are no data with which to compare the pharmacokinetic properties of the extended-release formulation and the suspension formulation. Until further data are available, the pharmacokinetic equivalence of the extended-release tablet and the suspension formulations should NOT be assumed.

Absorption and Metabolism

The extent of absorption (AUC) and the maximum plasma concentration (C_{max}) of cefaclor from cefaclor extended-release tablets are greater when the extended release tablet is taken with food.

[NOTE: The extent of absorption (AUC) of cefaclor from cefaclor immediate-release capsules is unaffected by food intake; however, when cefaclor immediate-release capsules are taken with food, the C_{max} is decreased.]

There is no evidence of metabolism of cefaclor in humans.

Comparative Serum Pharmacokinetics

Serum pharmacokinetic parameters for cefaclor extended-release tablets and cefaclor immediate-release capsules are shown in the table below.

TABLE 1
Comparative Pharmacokinetics of
Cefaclor Immediate-Release Capsules vs. Cefaclor Extended-Release Tablets
In Fasting and Fed States

Parameter	Cefaclor Extended-Release Tablets 375 mg		Cefaclor Extended-Release Tablets 500 mg		Cefaclor Immediate-Release Capsules 2 x 250 mg	
	fed n=10	fast NA	fed n=16	fast n=16	fed n=15	fast n=16
C_{max}	3.7 (1.1)	NA	8.2 (4.2)	5.4 (1.6)	9.3 (2.7)	16.8 (4.7)
T_{max}	2.7 (1.0)	NA	2.5 (0.8)	1.5 (0.7)	1.5 (0.6)	0.9 (0.4)
AUC	9.9 (2.2)	NA	18.1 (4.2)	14.8 (4.0)	20.5 (2.8)	19.2 (5.0)

(± 1 standard deviation)

NA = data not available

No drug accumulation was noted when cefaclor extended-release was given twice daily.

The plasma half-life in healthy subjects is independent of dosage form and averages approximately 1 hour.

Food Effect on Pharmacokinetics

When cefaclor extended-release tablet is taken with food, the AUC is 10% lower while the C_{max} is 12% lower and occurs 1 hour later compared to cefaclor immediate-release capsules. In contrast, when cefaclor extended-release tablet is taken without food, the AUC is 23% lower while the C_{max} is 67% lower and occurs 0.6 hours later, using an equivalent milligram dose of cefaclor immediate-release capsules as a reference. **Therefore, cefaclor extended-release should be taken with food.**

Special Populations

Renal Insufficiency

In patients with reduced renal function, the serum half-life of cefaclor is slightly prolonged. In those with complete absence of renal function, the plasma half-life of the intact molecule is 2.3 to 2.8 hours. Excretion pathways in patients with markedly impaired renal function have not been determined. Hemodialysis shortens the half-life by 25% to 30%.

Geriatric Patients

In elderly subjects (over age 65) with normal serum creatinine values, higher peak plasma concentration and AUCs have been observed. This is considered to be primarily a result of an age-related decrement in renal function, and has no apparent clinical significance. Therefore, dosage adjustment is not necessary in elderly subjects with normal serum creatinine values.

Microbiology

Cefaclor has *in vitro* activity against a broad range of gram-positive and gram-negative bacteria. The bactericidal action of cefaclor results from inhibition of cell-wall synthesis. Cefaclor is stable in the presence of some bacterial β -lactamases; consequently, some β -lactamase-producing organisms may be susceptible to cefaclor. Cefaclor extended-release tablet has been shown to be active against most strains of the following microorganisms both *in vitro* and in clinical infections as described in the INDICATIONS AND USAGE section.

Gram-positive aerobes

Staphylococcus aureus
Streptococcus pneumoniae
Streptococcus pyogenes

NOTE: Cefaclor is inactive against methicillin-resistant staphylococci.

Gram-negative aerobes

Haemophilus influenzae (non- β -lactamase-producing strains only)

Moraxella catarrhalis (including β -lactamase-producing strains)

The following *in vitro* data are available, but their clinical significance is unknown. Cefaclor exhibits *in vitro* minimum inhibitory concentrations (MICs) of 8 mcg/mL or less (systemic susceptibility breakpoint) against most ($\geq 90\%$) strains of the following microorganisms; however, the safety and effectiveness of cefaclor extended-release in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled trials.

Gram-positive aerobes

Staphylococcus epidermidis

Gram-negative aerobes

Haemophilus parainfluenzae

Klebsiella pneumoniae

Anaerobic bacteria

Peptococcus niger

Peptostreptococci

Propionibacterium acnes

NOTE: *Acinetobacter calcoaceticus*, *Enterobacter* spp., *Enterococcus* spp., *organella morgani*, *Proteus vulgaris*, *Providencia* spp., *Pseudomonas* spp., and *Serratia* spp. are resistant to cefaclor.

Susceptibility Testing

Dilution Techniques

Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure. Standardized procedures are based on a dilution method¹ (broth, agar or microdilution) or equivalent with standardized inoculum concentrations and standardized amounts of cefaclor powder. The MIC values should be interpreted according to the following criteria:

MIC (mcg/mL)	Interpretation
≤ 8	Susceptible (S)
16	Intermediate (I)
≥ 32	Resistant (R)

A report of "susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound in blood reaches the concentrations usually achievable. A report of "intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "resistant" indicates that the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable; other therapy should be selected.

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard cefaclor powder should provide the following MIC values:

Microorganism	ATCC	MIC range (mcg/mL)
<i>E. coli</i>	ATCC 25922	1 - 4
<i>E. faecalis</i>	ATCC 29212	> 32
<i>S. aureus</i>	ATCC 29213	1 - 4
<i>H. influenzae</i>	ATCC 49766*	1 - 4

*Broth microdilution tests performed using Haemophilus Test Medium (HTM)¹

Diffusion Techniques

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure² requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 30-mcg cefaclor to test the susceptibility of microorganisms to cefaclor.

Reports from the laboratory providing results of the standard single-disk susceptibility test with a 30-mcg cefaclor disk should be interpreted according to the following criteria:

Zone diameter (mm)	Interpretation
≥ 18	Susceptible (S)
15 - 17	Intermediate (I)
≤ 14	Resistant (R)

When testing * *H. influenzae*, the following interpretive criteria should be used:

Zone diameter (mm)	Interpretation
≥ 20	Susceptible (S)
17 - 19	Intermediate (I)
≤ 16	Resistant (R)

*Disk susceptibility tests performed using Haemophilus Test Medium (HTM)²

Interpretation should be as stated above for results using dilution techniques. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for cefaclor.

As with standardized dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion technique, the 30-mcg cefaclor disk should provide the following zone diameters in these laboratory test quality control strains:

Microorganisms	ATCC	Zone Diameter (mm)
<i>E. coli</i>	ATCC 25922	23 - 27
<i>S. aureus</i>	ATCC 29213	27 - 31
<i>H. influenzae</i> *	ATCC 49766	25 - 31

*Disk susceptibility tests performed using Haemophilus Test Medium (HTM)²

INDICATIONS AND USAGE

The safety and effectiveness of cefaclor extended-release in treating some of the indications and pathogens for which other formulations of cefaclor are approved have NOT been established. When administered at the recommended dosages and durations of therapy, cefaclor extended-release tablets are indicated for the treatment of patients with the following mild to moderate infections when caused by susceptible strains of the designated organisms. (See DOSAGE AND ADMINISTRATION AND CLINICAL STUDIES)

Acute bacterial exacerbations of chronic bronchitis due to *Haemophilus influenzae* (non- β -lactamase-producing strains only), *Moraxella catarrhalis* (including β -lactamase-producing strains) or *Streptococcus pneumoniae*.

Note: In view of the insufficient numbers of isolates of β -lactamase-producing strains of *Haemophilus influenzae* that were obtained from clinical trials with cefaclor extended-release for patients with acute bacterial exacerbations of chronic bronchitis or secondary bacterial infections of acute bronchitis, it was not possible to adequately evaluate the effectiveness of cefaclor extended-release for bronchitis known, suspected, or considered potentially to be caused by β -lactamase-producing *H. influenzae*.

Secondary bacterial infections of acute bronchitis due to *Haemophilus influenzae* (non- β -lactamase-producing strains only), *Moraxella catarrhalis* (including β -lactamase-producing strains), or *Streptococcus pneumoniae*. (See above NOTE)

Pharyngitis and tonsillitis due to *Streptococcus pyogenes*.

NOTE: Only penicillin by the intramuscular route of administration has been shown to be effective in the prophylaxis of rheumatic fever. Cefaclor extended-release tablet is generally effective in the eradication of *S. pyogenes* from the oropharynx; however, data establishing the efficacy of cefaclor extended-release tablet for the prophylaxis of subsequent rheumatic fever are not available.

Uncomplicated skin and skin structure infections due to *Staphylococcus aureus* (methicillin-susceptible).

NOTE: In view of the insufficient numbers of isolates of *Streptococcus pyogenes* that were obtained from clinical trials with cefaclor extended-release tablets for patients with uncomplicated skin and skin structure infections, it was not possible to adequately evaluate the effectiveness of cefaclor extended-release tablets for skin infections known, suspected, or considered potentially to be caused by *S. pyogenes*.

CONTRAINDICATIONS

Cefaclor extended-release tablets are contraindicated in patients with known hypersensitivity to cefaclor and other cephalosporins.

WARNINGS

BEFORE THERAPY WITH CEFACTOR EXTENDED-RELEASE IS INSTITUTED, CAREFUL INQUIRY SHOULD BE MADE TO DETERMINE WHETHER THE PATIENT HAS HAD PREVIOUS HYPERSENSITIVITY REACTIONS TO CEFACTOR, CEPHALOSPORINS, PENICILLINS, OR OTHER DRUGS. IF THIS PRODUCT IS TO BE GIVEN TO PENICILLIN-SENSITIVE PATIENTS, CAUTION SHOULD BE EXERCISED BECAUSE CROSS-SENSITIVITY AMONG BETA-LACTAM ANTIBIOTICS HAS BEEN CLEARLY DOCUMENTED AND MAY OCCUR IN UP TO 10% OF PATIENTS WITH A HISTORY OF PENICILLIN ALLERGY. IF AN ALLERGIC REACTION TO CEFACTOR EXTENDED-RELEASE OCCURS, DISCONTINUE THE DRUG. SERIOUS ACUTE HYPERSENSITIVITY REACTIONS MAY REQUIRE TREATMENT WITH EPINEPHRINE AND OTHER EMERGENCY MEASURES, INCLUDING OXYGEN, INTRAVENOUS FLUIDS, INTRAVENOUS ANTIHISTAMINES, CORTICOSTEROIDS, PRESSOR AMINES, AND AIRWAY MANAGEMENT, AS CLINICALLY INDICATED.

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including cefaclor, and may range from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth by clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is a primary cause of "antibiotic-associated colitis".

After the diagnosis of pseudomembranous colitis has been established, therapeutic measure should be initiated. Mild cases of pseudomembranous colitis usually respond to discontinuation of the drug alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation and treatment with an antibacterial drug clinically effective against *Clostridium difficile*.

PRECAUTIONS

General

Superinfection (overgrowth by non-susceptible organisms) should always be considered a possibility in a patient being treated with a broad spectrum antimicrobial. Careful observation of the patient is essential. If superinfection occurs during therapy, appropriate measures should be taken.

Drug Interactions

Antacids

The extent of absorption of cefaclor extended-release tablet is diminished if magnesium or aluminum hydroxide-containing antacids are taken within 1 hour of administration; H₂ blockers do not alter either the rate or the extent of absorption of cefaclor extended-release tablets.

Probenecid

The renal excretion of cefaclor is inhibited by probenecid.

Warfarin

There have been rare reports of increased prothrombin time with or without clinical bleeding in patients receiving cefaclor and warfarin concomitantly. No specific studies have been performed to rule in or rule out this potential drug/drug interaction.

Laboratory Test Interactions

CEFACTOR EXTENDED-
RELEASE TABLETS USP

0204-02



0204-02

CEFACTOR EXTENDED-
RELEASE TABLETS USP

Administration of cefaclor extended-release tablets may result in a false-positive reaction for glucose in the urine. This phenomenon has been seen in patients taking cephalosporin antibiotics when the test is performed using Benedict's and Fehling's solutions and also with Clinistest® tablets.

Carcinogenesis, Mutagenesis, Impairment of Fertility
 Studies in animals have not been performed to evaluate the carcinogenic or mutagenic potential for cefaclor. Reproduction studies have revealed no evidence of impaired fertility.

Usage in Pregnancy

Teratogenic Effect

Pregnancy Category B

Reproduction studies using cefaclor have been performed in mice, rats and ferrets at doses up to 3-5 times the maximum human dose (1500 mg/day) based on mg/m². These studies have revealed no harm to the fetus due to cefaclor. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, cefaclor extended-release tablets should be used during pregnancy only if clearly needed.

Labor and Delivery

Cefaclor extended-release tablets have not been studied for use during labor and delivery. Treatment should be given only if clearly needed.

Nursing Mothers

No studies in lactating women have been performed with cefaclor extended-release tablets. Small amounts of cefaclor (≤ 0.21 mcg/mL) have been detected in human milk following administration of single 500-mg doses of cefaclor. The effect on nursing infants is not known. Caution should be exercised when cefaclor extended-release tablet is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of cefaclor extended-release tablets in pediatric patients less than 16 years of age have not been established.

Geriatric Use

Healthy geriatric volunteers (≥65 years old) who received a single 750-mg dose of cefaclor extended-release tablets had 40% - 50% higher AUC and 20% lower renal clearance values when compared to healthy adult volunteers less than 45 years of age. These differences are considered to be primarily a result of age-related decreases in renal function. In clinical studies when geriatric patients received the usual recommended adult doses, clinical efficacy and safety were comparable to results in non-geriatric adult patients. No dosage changes are recommended for healthy geriatric patients.

ADVERSE REACTIONS

Clinical Trials

There were 3272 patients treated with multiple doses of cefaclor extended-release tablets in controlled clinical trials and an additional 211 subjects in pharmacology studies. There were no deaths in these trials thought to be related to toxicity from cefaclor extended-release tablets. Treatment was discontinued in 1.7% of patients due to adverse events thought to be possibly or probably drug-related.

The following adverse clinical and laboratory events were reported during the cefaclor extended-release tablet clinical trials conducted in North America at doses of 375 mg or 500 mg BID; however, relatedness of the adverse events to the drug was not assigned by clinical investigators during the trials (see Tables 2 and 3).

Table 2
Adverse Clinical Events
Cefaclor Extended-Release Tablets Multiple-Dose Dosing Regimens
Clinical Trials - North America
(n=1400)

Event	Incidence
Headache	4.9%
Rhinitis	3.9%
Diarrhea	3.8%
Nausea	3.4%
Vaginitis*	2.4%
Vaginal Moniliasis*	2.2%
Abdominal Pain	1.6%
Cough Increased	1.5%
Pharyngitis	1.4%
Pruritus	1.4%
Back Pain	1.0%

* n=934 for these events (subset of female participants).

Adverse reactions occurring during the clinical trials with cefaclor extended-release tablets with an incidence of less than 1% but greater than 0.1% included the following (listed alphabetically):

Accidental injury, anorexia, anxiety, arthralgia, asthma, bronchitis, chest pain, chills, congestive heart failure, conjunctivitis, constipation, dizziness, dysmenorrhea, dyspepsia, dysuria, ear pain, edema, fever, flatulence, flu syndrome, gastritis, infection, insomnia, leukorrhea, lung disorder, maculopapular rash, malaise, menstrual disorder, myalgia, nausea and vomiting, neck pain, nervousness, nocturia, otitis media, pain, palpitation, peripheral edema, rash, respiratory disorder, sinusitis, somnolence, surgical procedure, sweating, tremor, urticaria, vomiting.

NOTE: One case of serum-sickness-like reaction was reported among the 3272 adult patients treated with cefaclor extended-release tablets during the controlled clinical trials. These reactions have also been reported with the use of cefaclor in other oral formulations and are seen more frequently in pediatric patients than in adults. These reactions are characterized by findings of erythema multiforme, rash, and other skin manifestations accompanied by arthritis/arthralgia, with or without fever, and differ from classic serum sickness in that there is infrequently associated lymphadenopathy and proteinuria, no circulating immune complexes and no evidence to date of sequelae of the reaction. While further investigation is ongoing, serum-sickness-like reactions appear to be due to hypersensitivity and more often occur during or following a second (or subsequent) course of therapy with cefaclor. Such reactions have been reported with overall occurrence ranging from 1 in 200 (0.5%) in one focused trial; to 2 in 8346 (0.024%) in overall clinical trials (with an incidence in pediatric patients in clinical trials of 0.055%); to 1 in 38,000 (0.003%) in spontaneous event reports. Signs and symptoms usually occur a few days after initiation of therapy and subside within a few days after cessation of therapy. Occasionally these reactions have resulted in hospitalization, usually of short duration (median hospitalization = 2 to 3 days, based on postmarketing surveillance studies). In those patients requiring hospitalization, the symptoms have ranged from mild to severe at the time of admission with more of the severe reactions occurring in pediatric patients.

Table 3
Adverse Clinical Laboratory Events
Cefaclor Extended-Release Tablets Multiple-Dose Dosing Regimens
Clinical Trials - North America

Event	Incidence
Albumin decreased	0.3%
Alkaline phosphatase increased	0.3%
ALT/SGPT increased	0.3%
Bilirubin total increased	0.3%
Blood urea nitrogen (BUN) increased	0.2%
Calcium decreased	0.7%
Creatine phosphokinase increased	0.7%
Creatinine increased	0.5%
Eosinophils increased	0.3%
Erythrocyte count decreased	0.3%
GGT increased	0.2%
Hemoglobin decreased	0.2%
Lymphocytes decreased	0.3%
Mean Cell Volume (MCV) increased	0.7%
Neutrophils segmented decreased	0.3%
Phosphorus increased	0.7%
Platelet count decreased	0.3%
Potassium increased	0.4%
Sodium decreased	0.3%
Sodium increased	0.4%

In Postmarketing Experience

In addition to the events reported during clinical trials with cefaclor extended-release tablets, the following adverse experiences are among those that have been reported during worldwide postmarketing surveillance: allergic reaction, anaphylactoid reaction, angioedema, face edema, hypotension, Stevens-Johnson syndrome, syncope, paresthesia, vasodilatation and vertigo.

Other Adverse Reactions Associated with Other Formulations of Cefaclor

In addition to the above, the following other adverse reactions and altered laboratory tests have been associated with cefaclor in other oral formulation:

Clinical

Severe hypersensitivity reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, and anaphylaxis, have been reported rarely. Anaphylactoid events may be manifested by solitary symptoms, including angioedema, edema (including face and limbs), paresthesias, syncope, or vasodilatation. Anaphylaxis may be more common in patients with a history of penicillin allergy. Rarely, hypersensitivity symptoms may persist for several months. Symptoms of pseudomembranous colitis may appear either during or after antibiotic treatment. (See **WARNINGS**).

Laboratory

Abnormal urinalysis, eosinophilia, leukopenia, neutropenia, transient elevations in AST, and transient thrombocytopenia have been reported.

Cephalosporin-Class Reactions

In addition to the adverse reactions listed above, the following adverse reactions and altered laboratory tests have been reported for cephalosporin-class antibiotics:

Clinical

Confusion, erythema multiforme, genital pruritus, hepatic dysfunction including cholestasis, hemolytic anemia, reversible hyperactivity, hypertonia, and reversible interstitial nephritis.

Laboratory

Positive direct Coombs' test.

OVERDOSAGE

The toxic symptoms following an overdose of cefaclor may include nausea, vomiting, epigastric distress, and diarrhea. The severity of the epigastric distress and the diarrhea are dose-related.

Absorption of drugs from the gastrointestinal tract may be decreased by giving activated charcoal, which, in many cases, is more effective than emesis or lavage. Consider charcoal instead of or in addition to gastric emptying. Repeated doses of charcoal over time may hasten elimination of some drugs that have been absorbed.

Although cefaclor is considered dialyzable, neither forced diuresis, peritoneal dialysis, hemodialysis, nor charcoal hemoperfusion have been demonstrated to be beneficial in an overdose of cefaclor.

DOSE AND ADMINISTRATION

The absorption of cefaclor extended-release tablet is enhanced when it is administered with food. (See **CLINICAL PHARMACOLOGY**). Therefore, cefaclor extended-release tablets should be administered with meals (i.e., at least within one hour of eating). The extended-release tablets should not be cut, crushed, or chewed.

See INDICATIONS AND USAGE for information about patients for whom cefaclor extended-release tablet is indicated.

NOTE: 500 mg BID of cefaclor extended-release tablet is clinically equivalent to 250 mg TID of cefaclor as an immediate-release capsule in those indications listed in the **INDICATIONS AND USAGE** section of this label. **500 mg BID of cefaclor extended-release tablet is NOT equivalent to 500 mg TID of other cefaclor formulations.**

Adults (age 16 years and older):

Type of Infection (as qualified in the INDICATIONS AND USAGE section of this labeling)	Total Daily Dose	Dose and Frequency	Duration
Acute Bacterial Exacerbations of Chronic Bronchitis due to <i>H. influenzae</i> (non-β-lactamase-producing strains only), <i>Moraxella catarrhalis</i> (including β-lactamase-producing strains), or <i>Streptococcus pneumoniae</i> (See INDICATIONS AND USAGE)	1000 mg	500 mg q 12 hours	7 days
Secondary Bacterial Infection of Acute Bronchitis due to <i>H. influenzae</i> (non-β-lactamase-producing strains only), <i>M. catarrhalis</i> (including β-lactamase-producing strains), or <i>S. pneumoniae</i> (See INDICATIONS AND USAGE)	1000mg	500 mg q 12 hours	7 days
Pharyngitis and/or tonsillitis due to <i>S. pyogenes</i>	750 mg	375 mg q 12 hours	10 days
Uncomplicated Skin and Skin Structure infections due to <i>S. aureus</i> (methicillin-susceptible strains) (See INDICATIONS AND USAGE)	750 mg	375 mg q 12 hours	7 - 10 days

Elderly patients with normal renal function do not require dosage adjustments.

HOW SUPPLIED

Cefaclor extended-release tablets USP are available as blue, unscored, oval-shaped tablets, debossed "X" and "4194" on one side and "500" on the other side containing cefaclor monohydrate equivalent to 500 mg anhydrous cefaclor, packaged in bottles of 100 and 500 tablets.

PHARMACIST: Dispense in a tight, light-resistant container as defined in the USP. Use child-resistant closure (as required).

SHOULD NOT BE CUT, CRUSHED OR CHEWED

Keep tightly closed.

Store at controlled room temperature 15°-30°C (59°-86°F) (See USP).

MANUFACTURED BY
 IVAX PHARMACEUTICALS, INC.
 MIAMI, FL 33137

0172
 01/03
 82

CEFACLOR EXTENDED-RELEASE TABLETS USP

CLINICAL STUDIES

ACUTE BACTERIAL EXACERBATIONS OF CHRONIC BRONCHITIS AND SECONDARY BACTERIAL INFECTIONS OF ACUTE BRONCHITIS:

In adequate and well-controlled clinical trials of cefaclor extended-release tablets in the treatment of acute bacterial exacerbations of chronic bronchitis (ABECB) and secondary bacterial infections of acute bronchitis (SBIAB), only 4 evaluable patients with ABECB and no evaluable patients with SBIAB had infections caused by β-lactamase-producing *H. influenzae*. Four patients do not provide adequate data upon which to judge clinical efficacy of cefaclor extended-release tablets against β-lactamase-producing *H. influenzae*.

UNCOMPLICATED SKIN AND SKIN STRUCTURE INFECTIONS

Cefaclor extended-release tablets (375 mg Q12H)(n=115) was compared to cefaclor immediate-release capsules (250 mg TID)(n=106) for the treatment of patients with uncomplicated skin and skin structure infections, including cellulitis, pyoderma, abscess and impetigo. Patients were treated for 7 to 10 days and were evaluated for clinical resolution and bacterial eradication approximately one week after completing therapy. To be evaluable, all patients had to have a recognized pathogen isolated from the skin infection just prior to the initiation of therapy. The results of this randomized, double-blinded, U.S. trial demonstrated:

- (1) overall clinical cure rates were 72% (83 of 115 patients) and 75% (80 of 106 patients), respectively, for cefaclor extended-release tablets and cefaclor immediate-release capsules [95% CI around the 3% difference = -16% to +9%],
- (2) overall bacteriologic eradication rates against *Staphylococcus aureus* were comparable (see Table 4).

Table 4
Clinical Response* in Patients With Skin and Skin Structure Infections

Outcome by Pathogen	Cefaclor Extended-Release Tablets	Cefaclor Immediate-Release Capsules
<i>Staphylococcus aureus</i>	67/95 (71%)	58/81 (71%)
<i>Streptococcus pyogenes</i>	10/16 (63%)	8/9 (89%)
Other streptococci	7/11 (64%)	5/6 (83%)
Total	84/122 (69%)	71/96 (74%)

*Cure plus improvement

REFERENCES

- National Committee for Clinical Laboratory Standards. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically-Third edition; Approved Standard NCCLS Document M7-A3, Vol 13, No 25, NCCLS, Villanova, PA, December 1993
- National Committee for Clinical Laboratory Standards. Performance Standards for Antimicrobial Disk Susceptibility Tests-Fifth edition; Approved Standard NCCLS Document M2-A5, Vol 13, No 24, NCCLS, Villanova, PA, December 1993

